

Nonlocal, noncommutative picture in quantum mechanics and distinguished canonical maps

T. Hakioglu

*Physics Department,
Bilkent University,
06533 Ankara, Turkey*

Classical nonlinear canonical (Poisson) maps have a distinguished role in quantum mechanics. They act unitarily on the quantum phase space and generate $\hbar$ -independent quantum canonical maps. It is shown that such maps act in the noncommutative phase space under the classical covariance. A crucial observation made is that under the classical covariance the local quantum mechanical picture can become nonlocal in the Hilbert space. This nonlocal picture is made equivalent by the Weyl map to a full noncommutative picture in the phase space formulation of the theory. The connection between the entanglement and nonlocality of the representation is explored and specific examples of the generation of entanglement by using the concept of generalized Bell states are provided. That the results have direct application in generating vacuum soliton configurations in the recently popular scalar field theories of noncommutative coordinates is also demonstrated.

PACS: 03.65-w (Quantum Mech.),
03.65.Sq (Semiclass. theories and appl's.),
03.65.Fd (Algebraic methods)

A. Introduction

Recently increased activity in noncommutative (field and gauge) theories¹ and the natural observation of noncommutative spatial coordinates in string physics² arose a flurry of interest in a certain nonstandard extension of the quantum mechanics within this noncommutative picture.³ In these theories the fundamental entity, the field, acquires an operator character by its dependence on the noncommutative coordinates (NCC). The NCC can arise in certain physical limits of a quantum theory. Recently, Bigatti and Susskind⁴ have suggested the quantum model in two space dimensions of a charged particle in the plane interacting with a perpendicular magnetic field in the limit the field strength goes to infinity to observe noncommutativity in the plane coordinates. Considering that the noncommutative gauge theories are related to the ordinary ones by certain transformations² it is reasonable to inquire whether more general results can be obtained on the nature of the interrelationship and the possibility of invertible transformations between the noncommutative and the ordinary formalisms. For instance, in the case of gauge theories the *ordinary* and the noncommutative theories are related by a *gauge equivalence preserving* map reminiscent of a contact transformation between the fields and the gauge parameters vice versa.² On the other hand, the *phase space* is spanned by the generalized coordinates and the formulation of quantum mechanics in the phase space explicitly has features of noncommutativity. This can be used as a natural testing ground to understand whether more general transformations can be found between the ordinary and noncommutative pictures. In our context here the noncommutative generalized coordinates $\hat{z} = (\hat{z}_1, \hat{z}_2)$ are the *generalized momentum* and *position* operators respectively satisfying $[\hat{z}_j, \hat{z}_k] = i\theta_{jk}$ where $j, k = (1, 2)$ and the noncommutativity parameter is $\theta_{jk} = \hbar J_{jk}$ with J_{jk} describing the symplectic matrix $\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$.

The equivalence of the one dimensional phase space to the two dimensional noncommutative geometry approach is already known. The generalization of the ideas presented here to noncommutative geometries may therefore be direct. For instance, in the case of Landau model this equivalence was shown in Ref. [5]. More recently it was proposed that there is an interesting connection between the quantum Hall effect and the noncommutative matrix models.^{6,7} The noncommutative picture can be invertibly converted into a bilocal coordinate picture by the Weyl map. Once a bilocal representation is obtained the (non)separability of these representations becomes a crucial property. When the BLC dependence is separable the corresponding noncommutative picture reduces to the standard (Schrödinger) picture. On the other hand, the nonseparability of the dependence on the bilocal coordinates requires a full phase space formalism (which may not be reducible to a pure coordinate or momentum

representation as in the standard Schrödinger picture). This full treatment of the noncommutative phase space and the corresponding nonlocal function space suggest an extended frame above the standard formulation of quantum mechanics. The purpose of this letter is to explore this extended quantum mechanical picture.

In section B the basic connections between the BLC and the phase space representations are established by the Weyl map and the question of the separability versus nonseparability in the BLC picture is studied. It is shown that the separable bilocal solutions are actually local and they can be fitted in the standard Schrödinger picture. The nonlocal solutions arise from the nonseparable bilocal coordinate dependence and they fit in the extended picture. In section C, it is shown that under the action of the standard linear Hilbert space operators the separable and the nonseparable functions are disjoint. In section D we examine the phase space approach in this extended quantum mechanical picture. By reaching beyond the standard Hilbert space operator methods, we find distinguished unitary (in the noncommutative -phase- space) isomorphisms (canonical maps) joining the separable and the nonseparable sectors. In the view of the continuous canonical transformations, these isomorphisms appear to be induced by the classical canonical (Poisson) maps. In section E a connection between the nonlocality and the entanglement is examined in the concept of *generalized Bell states* is introduced. Specific nonlinear Poisson maps are given in section E that produce these states from the pure states. Recently the generalized Bell states have been observed in the vacuum soliton solutions of the noncommutative field theories and this concept may be required in building a genuine equivalence scheme for the set of all noncommutative soliton solutions. This particular application is timely and relevant arising as an effect of the Poisson maps which is also discussed in a paragraph therein. The section F is a schematical illustration and summary of the results in the work.

B. The extended quantum mechanical picture

The ordinary quantum mechanics is the sector of the field theory supporting a single degree of freedom. We adopt an operator-like approach in this description^{8,9} that will be referred to as the *waveoperator*. The waveoperator is a complex functional of the noncommutative generalized coordinates which will be denoted by $\hat{\rho} = \rho(\hat{z})$. In this representation quantum mechanics is formulated by extremizing the action⁸

$$S = \int dt \text{Tr} \left\{ i\hbar \hat{\rho}^\dagger \partial_t \hat{\rho} - \hat{\rho}^\dagger \hat{\mathcal{H}} \hat{\rho} \right\}$$

$$\hat{\mathcal{H}} = \mathcal{H}(\hat{z}_1, \hat{z}_2) = \frac{\hat{z}_1^2}{2m} + V(\hat{z}_2) \quad (1)$$

with respect to some coordinate space description $\rho(y, x) = \langle y | \hat{\rho} | x \rangle$. Here $\hat{\mathcal{H}}$ is a Hamiltonian operator in

the Hilbert space and Tr stands for the trace in the position basis $|x\rangle$ (i.e. $\hat{z}_2|x\rangle = x|x\rangle$). Throughout the paper, the waveoperator is time dependent although we will not write it explicitly.

Evaluating the action by considering the trace in this position basis, and minimizing it with respect to $\rho^*(x, y) = \langle y|\hat{\rho}|x\rangle$ we obtain

$$i\hbar \partial_t \rho(y, x) = \int du h(y, u) \rho(u, x) \quad (2)$$

where $\rho(y, x)$ is the representation of the waveoperator in the *bilocal coordinates* (BLC) y, x . Equivalently the bilocal $\rho(y, x)$ can be mapped isomorphically onto a doubled Hilbert space $H \times H$ and this procedure is known as the Gel'fand-Naimark-Segal construction. The $H \times H$ representations have been noticed recently to be of use in the phase space representations of the quantum canonical Lie algebras¹¹. In this scheme $\rho(y, x)$ can be referred as the wavefunction in this doubled Hilbert space¹². Similarly, $h(y, u) = \langle y|\mathcal{H}|u\rangle$ is the Hamiltonian in BLC. Eq. (2) will be referred as the bilocal Schrödinger equation.

Note that, (2) does not have unique solutions. The nonuniqueness arises from the indeterminacy in the x dependence. In order to show this indeterminacy in (2) we assume that one solution is separable, e.g. $\rho(y, x) = \psi_1(y) \psi_2(x)$ where both ψ_k , $k = 1, 2$ have finite norm¹⁰. Here $\psi_2(x)$ must be time independent. Inserting this in (2) one obtains the standard Schrödinger equation for ψ_1 whereas since ψ_2 is time independent, it appears as a multiplicative function on both sides and thus it cannot be determined by the equation. In order to reach beyond this standard representation the nonseparable solutions of (2) in terms of the BLC, e.g. $\rho(y, x) \neq \psi_1(y) \psi_2(x)$ are of interest.

The separable and nonseparable sectors in the space of bilocal functions can be connected to each other by certain types of canonical maps. This can be best understood if we use the Weyl map¹³ to transform $\rho(y, x)$ into a function $\tilde{\rho}(z)$ where $z = (z_1, z_2)$ are the generalized coordinates in some noncommutative space $\mathcal{Z}_*$. Note that throughout the paper a generic dependence on (z_1, z_2) will be denoted by z and functions of z will be hatted with a tilde. Also particularly in this work, z_1, z_2 describe the phase space coordinates, e.g. the generalized canonical coordinate and momentum respectively. The standard approach to the phase space quantum mechanics is to use the Weyl correspondence which is an analytic and invertible map from an arbitrary Hilbert space operator $\hat{O}$ to a function $\tilde{o}(z)$ in $\mathcal{Z}_*$. This correspondence will be denoted by $\mathcal{W} : \hat{O} \leftrightarrow \tilde{o}(z)$. Since a Hilbert space operator can also be represented by a bilocal function as $\hat{O} \leftrightarrow \langle y|\hat{O}|x\rangle = o(y, x)$, the combination of these two is also a well-defined (Weyl) map $W : o(y, x) \leftrightarrow \tilde{o}(z)$. The Weyl map is explicitly given by

$$o(y, x) = \int \frac{d^2 z}{2\pi\hbar} \mathcal{K}_z(y, x) \tilde{o}(z). \quad (3)$$

Here $\mathcal{K}_z(y, x) = e^{iz_1(x-y)/\hbar} \delta(z_2 - \frac{x+y}{2})$ is the invertible Weyl kernel. The function $\tilde{o}(z)$ is the standard Weyl symbol of $\hat{O}$ in the phase space $\mathcal{Z}_*$. In this context, o and $\tilde{o}$ are two different representations of the Hilbert space operator $\hat{O}$. We therefore have the following triangle diagram

$$\begin{array}{ccc} \hat{O} & & \tilde{o}(z) \\ & \searrow & \nearrow \\ & o(y, x) & \end{array} \quad \begin{array}{c} \xleftarrow{\mathcal{W}} \\ \\ \xleftarrow{W} \end{array} \quad (4)$$

Note that W is a well defined map by itself between a bilocal function $o(y, x)$ and the function $\tilde{o}(z)$. It can exist as a transformation independently from $\mathcal{W}$, a fact which we exploit later in sections D.3 and D.5.

The noncommutativity of the coordinates in $\mathcal{Z}_*$ is encoded in the associative $\star$ -product which is implicit in (3). The $\star$ -product is best described by the Weyl map $W : [\int du o_1(y, u) o_2(u, x)] \leftrightarrow \tilde{o}_1(z) \star \tilde{o}_2(z)$ where

$$\begin{aligned} \tilde{o}_1 \star \tilde{o}_2 &= \tilde{o}_1 \exp\left\{\frac{i}{2} \overleftarrow{\partial}_{z_i} \theta_{ij} \overrightarrow{\partial}_{z_j}\right\} \tilde{o}_2 \\ &\neq \tilde{o}_2 \star \tilde{o}_1 \end{aligned} \quad (5)$$

with the arrows indicating the direction that the partial derivatives act. The measure of the noncommutativity of $\tilde{o}_1$ and $\tilde{o}_2$ in $\mathcal{Z}_*$ is provided by the Moyal bracket (MB) $\{\tilde{o}_1, \tilde{o}_2\}_z^{(M)} = \tilde{o}_1 \star \tilde{o}_2 - \tilde{o}_2 \star \tilde{o}_1$. We now consider for $\hat{O}$ the waveoperator $\hat{\rho}$. Then, $o(y, x)$ and $\tilde{o}(z)$ correspond to $\rho(y, x)$ and $\tilde{\rho}(z)$ respectively. Using (3) the Weyl map of (2) is found to be

$$i\hbar \partial_t \tilde{\rho}(z) = \tilde{h}(z) \star \tilde{\rho}(z). \quad (6)$$

Here $\tilde{h}$ is the Weyl map of the bilocal Hamiltonian h in (2). We refer to Eq. (6) as the $\star$ -Schrödinger equation which is basically an operator relation manifested by the waveoperator interpretation [Note that if $\hat{\rho}$ described the standard quantum mechanical density operator, the right hand side of (6) would not be given by a single $\star$ -product but by a MB instead. Hence (6) would be the Moyal equation for the density operator $i\hbar \partial_t \hat{\rho} = \tilde{h} \star \hat{\rho} - \hat{\rho} \star \tilde{h} = \{\tilde{h}, \hat{\rho}\}^{(M)}$].

If $\tilde{\rho}$ is an existing solution of (6), a new solution can be obtained by $\star$ -multiplying that equation from the right by *time independent* function $\tilde{\xi}(z)$ such that $-i\hbar \partial_t \tilde{\xi}(z) = \tilde{\xi}(z) \star \tilde{h} = 0$. The new solution is then represented by $\tilde{\rho}'(z) = \tilde{\rho} \star \tilde{\xi}$ and respects $i\hbar \partial_t \tilde{\rho}' = \{\tilde{h}, \tilde{\rho}'\}^{(M)}$ with unique solutions (upto initial conditions).

In the separable case, Eq. (6) has also a specific physical realization in terms of the generalized Wigner function. Consider the case $\hat{\rho} = |\phi_E\rangle \langle \chi|$ where E labels some basis states $|\phi_E\rangle$ which are the eigen solutions of the Schrödinger equation with some Hamiltonian and $|\chi\rangle$ is a time independent state under the same Hamiltonian. In this separable form $\rho(y, x) = \phi_E(y) \chi^*(x)$ is an eigen solution of (2) and the $\tilde{\rho}(z)$ is the corresponding eigen

solution of (6) which, by the application of (3), is given by

$$\tilde{\rho}(z) = \int dx e^{-i z_1 x/\hbar} \phi_E(z_2 - \frac{x}{2}) \chi^*(z_2 + \frac{x}{2}). \quad (7)$$

which is the generalized Wigner function¹⁴ W_{χ, ϕ_E} . If the basis states $|\phi_E\rangle$ are the energy eigen basis of the Hamiltonian $\hat{\mathcal{H}}$ with energy E then (7) is the eigen solution of Eq. (6).¹⁵

Here $\rho(y, x) = \phi_E(y) \chi^*(x)$ is manifestly separable by choice, where the local solution is indicated by $\phi_E(y)$. In order to explore beyond the ordinary local sector we must find how to reach the nonseparable sector of the doubled Hilbert space.

C. Separability and the Hilbert space operators

We look for connections between the local $H \times H$ subsector and the general doubled Hilbert space. Hence the relevant question is changing the separability of the functions like $\rho(y, x)$. Within the frame of standard transformations in quantum mechanics the answer is simple. Consider the Hilbert space operator -unitary or nonunitary- (speaking of the Hilbert space the standard -local- quantum mechanical Hilbert space is implied and not the doubled Hilbert space) $\hat{\Omega}$ acting on $\hat{\rho}$ as

$$\hat{\Omega} : \hat{\rho} = \hat{\rho}' = \hat{\Omega} \hat{\rho} \hat{\Omega}^\dagger \quad (8)$$

Suppose that we start from a separable case in (8) as $\hat{\rho} = |\psi_1\rangle\langle\psi_2|$, e.g. $\rho(y, x) = \psi_1(y) \psi_2(x)$. Then $\hat{\rho}' = \hat{\Omega} |\psi_1\rangle\langle\psi_2| \hat{\Omega}^\dagger = |\psi'_1\rangle\langle\psi'_2|$, where $\|\psi_j\| = \|\psi'_j\|$. We observe that $\hat{\rho}'$ is still separable, e.g. $\rho'(y, x) = \psi'_1(y) \psi'_2(x)$. Likewise, the invertibility of $\hat{\Omega}$ ensures that, the nonseparable $\rho(y, x)$ is transformed into a nonseparable $\rho'(y, x)$. Therefore Eq. (8) is a simple proof that the separable and the nonseparable BLC representations are disjoint within the reach of standard transformation $\hat{\Omega}$. Our discussion here should give us the clue that in reaching for the nonseparable solutions one needs to go beyond $H \times H$. In searching for the answer, an extended view of the unitary (canonical) transformations in $\mathcal{Z}_\star$ will be given in the following section.

D. An extended view of quantum canonical maps

One alternative way to look into the unitary transformations is by canonical maps in $\mathcal{Z}_\star$. Consider $\hat{z}_j$ and $\hat{Z}_j$ as the old and the new noncommutative generalized coordinates. A canonical map, in the operator picture, is given by $\hat{\Omega} : \hat{z}_j \mapsto \hat{Z}_j = Z_j(\hat{z}_1, \hat{z}_2)$, ($j = 1, 2$) such that the canonical commutation relations are preserved

$$\hat{\Omega} : [\hat{z}_j, \hat{z}_k] = [\hat{Z}_j, \hat{Z}_k] = i \theta_{jk}. \quad (9)$$

The corresponding relations in $\mathcal{Z}_\star$ can be obtained by the Weyl transform and are based on the *canonical* MB, where the latter is defined by $\{z_i, z_j\}^{(M)} = i \theta_{ij} = \{Z_i, Z_j\}^{(M)}$. The *canonical* MB is *basis independent* [see Eq. (21) below] whereas a general MB is not. Because of this, we will choose the z basis in expressing the $\star$ products and MBs, hence $\star = \star_z$ as given by (5). Here $Z_j = Z_j(z_1, z_2)$, $j = 1, 2$ are the Weyl symbols of the new noncommutative generalized coordinates $\hat{Z}_j$. In the following two different types of canonical maps will be examined as classical and quantum. These two types are distinguished by their covariance properties when they act on the functions of z . We refer to Ref. [16] for a detailed comparative study of the classical and the quantum cases.

1. Classically covariant canonical maps $\Omega_{\mathcal{Z}}$: type-I

Since we are dealing with the classical case, a necessity now arises to differentiate the classical commutative space $\mathcal{Z}$ from the noncommutative one $\mathcal{Z}_\star$. The commutativity in $\mathcal{Z}$ is induced by the product $\tilde{o}_1(z) \tilde{o}_2(z) = \tilde{o}_2(z) \tilde{o}_1(z)$ as opposed to the noncommutative one in $\mathcal{Z}_\star$ as expressed in (5). However there are also common features. For instance, a square integrable function in $\mathcal{Z}_\star$ is also square integrable in $\mathcal{Z}$ due to the integral property of the $\star$ -product $\int d^2 z \tilde{o}_1(z) \star_z \tilde{o}_2(z) = \int d^2 z \tilde{o}_1(z) \tilde{o}_2(z)$. This allows us to refer to the functions of z without referring to the underlying commutative or noncommutative space. The representations of the operators, however, generally differ due to the different covariance properties of them in $\mathcal{Z}$ or $\mathcal{Z}_\star$.

With this in mind, we assume that an infinitesimal generator $\mathcal{G}(z)$ of the canonical map exists generating the first order infinitesimal changes in the function $\tilde{o}(z)$, i.e. $\tilde{o} \rightarrow \tilde{o} + \delta \tilde{o}$, by the Poisson Bracket (PB). This is given by the classical textbook formula, $\delta \tilde{o}(z) = \epsilon \{\mathcal{G}, \tilde{o}\}^{(P)}$ where a general PB is given between two such functions $\tilde{o}_1$ and $\tilde{o}_2$ by $\{\tilde{o}_1, \tilde{o}_2\}^{(P)} \equiv J_{jk} (\partial_{z_j} \tilde{o}_1) (\partial_{z_k} \tilde{o}_2)$. According to the standard phase space analytical mechanics, $\mathcal{G}$ is transformed into a Hamiltonian vector field as $X_{\mathcal{G}} = J_{jk} (\partial_{z_j} \mathcal{G}) (\partial_{z_k})$ which generates the same infinitesimal change by the Lie bracket $\delta \tilde{o}(z) = \epsilon [X_{\mathcal{G}}, \tilde{o}] = \epsilon (X_{\mathcal{G}} \tilde{o} - \tilde{o} X_{\mathcal{G}})$. The classical canonical maps $\Omega_{\mathcal{Z}}$ we consider here are finite transformations in $\mathcal{Z}$ obtained by exponentiating the classical Hamiltonian vector fields as

$$\Omega_{\mathcal{Z}} = e^{\epsilon X_{\mathcal{G}}} \quad (10)$$

The finite action of $\Omega_{\mathcal{Z}}$ on the functions of z is given by

$$\begin{aligned} \Omega_{\mathcal{Z}} : \tilde{o} = \tilde{o} + \epsilon [X_{\mathcal{G}}, \tilde{o}] + \dots \\ + \frac{\epsilon^n}{n!} \underbrace{[X_{\mathcal{G}}, \dots [X_{\mathcal{G}}, [\tilde{o}, X_{\mathcal{G}}] \dots]}_n + \dots \end{aligned} \quad (11)$$

which can be shown to have the manifest property

$$\Omega_{\mathcal{Z}} : \tilde{o}(z) = \tilde{o}(Z) \quad (12)$$

where $Z = \Omega_{\mathcal{Z}} : z$. Another way of writing (12) is

$$\begin{aligned} [\Omega_{\mathcal{Z}} : \tilde{o}_1(z) \tilde{o}_2(z)] &= \tilde{o}'_1(z) \tilde{o}'_2(z) \\ &= \tilde{o}_1(Z) \tilde{o}_2(Z) . \end{aligned} \quad (13)$$

which will be a useful relation to facilitate the comparison with the quantum case. Eq's (10-13) are well-known results that can be found in the books on Lie algebraic techniques in analytical mechanics.¹⁷ In (13) the square brackets indicate that there are no other operators to the left or right acted upon by the expressions inside and the primes denote the transformed functions of z . This notation will be used throughout the paper. We note that Eq. (13) defines the *classical covariance* as used in the context of this work.¹⁶ The action of the canonical map $\Omega_{\mathcal{Z}}$ on the Poisson Bracket (PB) can also be defined similarly. Denoting the latter by $\{\tilde{o}_1, \tilde{o}_2\}^{(P)} \equiv J_{jk}(\partial_{z_j} \tilde{o}_1)(\partial_{z_k} \tilde{o}_2)$, the classical covariance implies

$$\begin{aligned} [\Omega_{\mathcal{Z}} : \{\tilde{o}_1(z), \tilde{o}_2(z)\}^{(P)}] \\ = \{[\Omega_{\mathcal{Z}} : \tilde{o}_1], [\Omega_{\mathcal{Z}} : \tilde{o}_2]\}^{(P)} \\ = \{\tilde{o}_1(Z), \tilde{o}_2(Z)\}^{(P)} . \end{aligned} \quad (14)$$

2. $\star$ -covariant canonical maps $\Omega_{\mathcal{Z}_\star}$: type-II

Consider a unitary operator $\hat{\Omega} = e^{i\epsilon \hat{G}}$ with $\epsilon \in \mathbb{R}$ and $\hat{G}$ Hermitian. $\hat{\Omega}$ acts on the operator $\hat{O}$ as

$$\begin{aligned} \hat{\Omega} : \hat{O} &= \hat{\Omega} \hat{O} \hat{\Omega}^\dagger \\ &= \hat{O} + i\epsilon [\hat{G}, \hat{O}] + \dots \\ &+ \frac{(i\epsilon)^n}{n!} \underbrace{[\hat{G} \dots [\hat{G}, [\hat{O}, \hat{G}] \dots]]}_n + \dots \end{aligned} \quad (15)$$

In analogy with the classical case, $\hat{G}$ above is the generator of the quantum canonical maps in the Hilbert space. We describe the Weyl symbol of $\hat{\Omega}$ by $\omega_{\mathcal{Z}_\star}(z)$ and its adjoint $\hat{\Omega}^\dagger$ by $\omega_{\mathcal{Z}_\star}^*(z)$. Such a map is given by the commuting diagram

$$\begin{array}{ccc} \hat{z}_j & \xleftarrow{\hat{\Omega}} & \hat{Z}_j \\ \mathcal{W} \downarrow & & \downarrow \mathcal{W} \\ z_j & \xleftarrow{\omega_{\mathcal{Z}_\star}} & Z_j \end{array} \quad \begin{array}{l} = \hat{\Omega} \hat{z}_j \hat{\Omega}^\dagger \\ \\ = \omega_{\mathcal{Z}_\star} : z_j \end{array} \quad (16)$$

It was shown^{16,18} that $\omega_{\mathcal{Z}_\star}$ acts on functions of z as

$$\omega_{\mathcal{Z}_\star} : \tilde{o}(z) = \tilde{o}'(z) = \omega_{\mathcal{Z}_\star}(z) \star_z \tilde{o}(z) \star_z \omega_{\mathcal{Z}_\star}^*(z) . \quad (17)$$

The canonical map $\omega_{\mathcal{Z}_\star}$ is the quantum analog of the classical map of $\Omega_{\mathcal{Z}}$. We actually gained an advantage in the search for the methods to reach beyond the Hilbert space operator approaches by formulating the canonical maps

in $\mathcal{Z}_\star$. The reason is that the quantum canonical maps can, in general, be handled in the phase space $\mathcal{Z}_\star$ in a similar way to the classical case in $\mathcal{Z}$ and this can be done in a totally independent way from the Hilbert space operator methods. Namely, the operator connections induced by the Weyl correspondence $\mathcal{W}$ can be totally and consistently ignored in (16). Examining (4), what remains is the Weyl map W between double (bilocal) Hilbert space and the phase space. We now proceed to discuss the $\star$ -covariance in Eq. (17).

If we use two such functions $\tilde{o}_1$ and $\tilde{o}_2$, Eq. (17) implies that

$$[\omega_{\mathcal{Z}_\star} : \tilde{o}_1 \star_z \tilde{o}_2] = [\omega_{\mathcal{Z}_\star} : \tilde{o}_1] \star_z [\omega_{\mathcal{Z}_\star} : \tilde{o}_2] . \quad (18)$$

Eq. (18) is to be regarded as the extended version of the classical covariance in (13). In the context of this work it will be referred to as the $\star$ -covariance¹⁶. Also note that, if we denote the canonical map in $\mathcal{Z}_\star$ by $\omega_{\mathcal{Z}_\star} : z_j \rightarrow Z_j$, Eq. (18) implies violation of the classical covariance, e.g. $[\omega_{\mathcal{Z}_\star} : \tilde{o}](z) \neq \tilde{o}(Z)$. Comparing (13) and (18) we note that $\Omega_{\mathcal{Z}}$ in (13) preserves the commutativity of the standard product between the functions $\tilde{o}_1$ and $\tilde{o}_2$ whereas for $\Omega_{\mathcal{Z}_\star}$ in (18) the noncommutativity by $\star$ -product is invariant. The MB is therefore transformed by $\omega_{\mathcal{Z}_\star}$ as

$$\begin{aligned} [\omega_{\mathcal{Z}_\star} : \{\tilde{o}_1(z), \tilde{o}_2(z)\}^{(M)}] \\ = \{[\omega_{\mathcal{Z}_\star} : \tilde{o}_1], [\omega_{\mathcal{Z}_\star} : \tilde{o}_2]\}^{(M)} \\ \neq \{\tilde{o}_1(Z), \tilde{o}_2(Z)\}^{(M)} . \end{aligned} \quad (19)$$

At the limit $\hbar \rightarrow 0$ the classical and star covariances coincide.

3. $\hbar$ -independent canonical maps:

A particular subgroup of canonical maps in types I and II is distinguished by its simultaneous representations both in $\mathcal{Z}$ and $\mathcal{Z}_\star$. This is the subgroup of $\hbar$ -independent canonical maps and may be crucial for a unified understanding of the classical and quantum phase spaces.

In what follows, we pay specific attention to this subgroup and derive some of its properties. Consider a canonically conjugated pair $Z_j = Z_j(z_1, z_2)$; ($j = 1, 2$) with $\{z_j, z_k\}^{(M)} = \{Z_j, Z_k\}^{(M)} = i\theta_{jk}$ where both the old z_k and the new Z_k canonical variables are assumed to be independent of $\hbar$. Expanding the MB in $\theta_{jk} = \hbar J_{jk}$ we have

$$\{Z_j, Z_k\}_z^{(M)} = i\hbar \{Z_j, Z_k\}_z^{(P)} + \mathcal{O}(\theta^3) \quad (20)$$

where the first term is the Poisson Bracket (PB) $\{Z_j, Z_k\}_z^{(P)} \equiv J_{\ell m}(\partial_{z_\ell} Z_j)(\partial_{z_m} Z_k)$. Since we assumed that Z_j 's have no dependence on $\hbar$, by equating (20) to $i\theta_{jk}$ and matching the powers of θ we deduce that the $\mathcal{O}(\theta^3)$ and higher order terms in (20) must all vanish. The nonvanishing part of (20) is

$$\begin{aligned}\{Z_j, Z_k\}_z^{(M)} &= i\hbar \{Z_j, Z_k\}_z^{(P)} \\ &= i\hbar J_{jk}\end{aligned}\quad (21)$$

which means that $\{Z_j, Z_k\}_z^{(P)} = J_{jk}$. This is the condition for the pair Z_1, Z_2 to be classically canonical. We learn that any $\hbar$ independent quantum canonical map $\omega_{Z_*} : z_j \rightarrow Z_j$ in (20) implies (21). Therefore such maps are also canonical in the Poisson sense and conversely all $\hbar$ -independent Poisson maps are also quantum canonical. We will refer to them as type-I as well. The corollary of this result is that an $\hbar$ independent quantum canonical pair can also be obtained by an appropriate Ω_Z . Now, we consider a second $\hbar$ -independent map Ω'_Z . By definition it acts classically on the canonical MB as

$$\begin{aligned}\Omega'_Z : \frac{1}{i\hbar} \{Z_j, Z_k\}^{(M)} &= \Omega'_Z : \{Z_j, Z_k\}^{(P)} \\ &= \{[\Omega'_Z : Z_j], [\Omega'_Z : Z_k]\}^{(P)} \\ &= \{Z_j(z'), Z_k(z')\}^{(P)} \\ &= J_{jk}\end{aligned}\quad (22)$$

where $z' = \Omega'_Z : z$. In (22) the second line is obtained by the classical covariance of the Poisson bracket in (14). The third line is an application of (13). The last line is the statement of the invariance of the canonical MB under $\hbar$ -independent canonical maps which is a corollary of (21). Therefore the classically covariant [see (13)] canonical maps define an $\hbar$ -independent automorphism on the *canonical* MB. That this result is correct only for canonical pairs and not for arbitrary functions of z is the essence of the $\star$ -covariance. In the literature, it almost goes without saying that all quantum canonical maps fall in the type-II. As we have seen above, type-I maps also preserve the canonical MB although their action is truly different from that of type-II. Therefore the space of canonical maps in Z_* is actually a union of types I and II which is referred to as the extended quantum phase space.

4. Nonseparability and the type-I maps

In the standard quantum mechanics, the canonical transformations of type-II have integral transforms. Denoting by $\phi(x)$ some local Hilbert space function, a typical map $\hat{\Omega}$ on $\phi(x)$ is expressed by

$$(\hat{\Omega} : \phi)(x') = \int dx' u(x, x') \phi(x') . \quad (23)$$

In the rest of this letter, we will be interested in examining similar integral transforms adopted for the bilocal representations and for canonical maps of type-I. For separable cases these bilocal transforms reduce to the direct products of the integral transforms like in (23) of which an example is given below from the linear canonical group. Denoting by Ω_Z a generic type-I map, its action can be written as

$$\begin{aligned}o'(y, x) &= (\Omega_Z : o)(y, x) \\ &= \int \frac{d^2 z}{2\pi\hbar} \mathcal{K}_z(y, x) \tilde{o}'(z)\end{aligned}\quad (24)$$

with $\tilde{o}'(z) = (\Omega_Z : \tilde{o})(z) = \tilde{o}(Z)$ as dictated by (13). Using the inverse of (3) we convert (24) into an integral transform between the old and the new functions of BLC as

$$o'(y, x) = \int du \int dv \mathcal{L}_{\Omega_Z}(y, x; v, u) o(v, u) \quad (25)$$

with the integral kernel

$$\mathcal{L}_{\Omega_Z}(y, x; u, v) = \int \frac{d^2 z}{2\pi\hbar} \mathcal{K}_z(y, x) (\Omega_Z : \mathcal{K}_z)(v, u) . \quad (26)$$

In Eq. (26) the classical covariance in (13) implies that $\Omega_Z : \mathcal{K}_z = \mathcal{K}_{\Omega_Z : z} = \mathcal{K}_Z$. Eq. (25) is the bilocal extension of (23). We now examine the separability of the general canonical map in (25) by three general examples.

a) Linear canonical group $\Omega_Z^{(a)} = \Lambda_g \in Sp_2(\mathbb{R})$ acts on the phase space as

$$\Lambda_g : \tilde{o}(z) = \tilde{o}(g^{-1} : z) , \quad g = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \quad (27)$$

where $\det g = 1$ and $g^{-1} : z = Z$ is the transformed coordinate. It is known that the linear canonical group is the only group of transformations for which the classical (type-I) and $\star$ -covariances (type-II) coincide. Based on this fact we already expect (27) not to have an effect on the separability. Nevertheless, the explicit calculation is illustrative. Using the inverse of the Weyl kernel in (3) we map the transformed and initial solutions by calculating the kernel in (26) for Λ_g as

$$\begin{aligned}\mathcal{L}_g &= \int \frac{d^2 z}{2\pi\hbar} \mathcal{K}_z(y, x) (\Lambda_g : \mathcal{K}_z)(v, u) \\ &= \int \frac{d^2 z}{2\pi\hbar} \mathcal{K}_z(y, x) \mathcal{K}_{g^{-1}:z}(v, u) \\ &= \exp\left\{\frac{i}{2\hbar c} [d(u^2 - v^2) + a(x^2 - y^2) - 2(xu - yv)]\right\}\end{aligned}\quad (28)$$

Here $\mathcal{L}_g$ is the kernel of the map Λ_g . As expected it is manifestly separable, i.e. $\mathcal{L}_g(y, x; u, v) = \mathcal{L}_g(0, x; u, 0) \mathcal{L}_g(y, 0; 0, v)$. Therefore the transformed solution is separable when $o(y, x) = \psi_1(y) \psi_2(x)$ and non-separable when $o(y, x)$ is nonseparable. In the first case both dependences on the coordinates are transformed separately and identically as

$$\begin{aligned}Y'(y) X'(x) &= \left[\int dv \mathcal{L}_g(y; v) Y(v) \right] \\ &\times \left[\int dv \mathcal{L}_g(x; u) X(u) \right] .\end{aligned}\quad (30)$$

where each square bracket is an integral transform of type-II as in (23).

b) As the second kind of type-I maps we consider

$$\Omega_{\mathcal{Z}}^{(b)} : \tilde{o}(z) = \tilde{o}(z_1 + \tau_1 A_1(z_2), z_2 + \tau_2 A_2(z_1)) \quad (31)$$

where it is crucial that the functions A_j 's are $\hbar$ independent. The free and real parameter τ_j is either momentum, ($\tau_1 = \tau, \tau_2 = 0$) or coordinate-like ($\tau_1 = 0, \tau_2 = \tau$). Two or higher dimensional versions of (31) are normally referred to as the gauge transformations (not considered here). Here we consider a momentum-like map ($\tau_1 = \tau \neq 0, \tau_2 = 0$). Using (31) in (3) one finds

$$o'(y, x) = e^{i\frac{\tau}{\hbar} x_- A(x_+)} o(y, x) \quad (32)$$

where $x_- = (x - y)$ and $x_+ = (x + y)/2$. Due to the exponential factor the transformation kernel in (32) is, for general cases, manifestly nonseparable. Exception is when $A(x_+) \propto \alpha + \beta x_+$, with α, β constant, in which case the exponential separates.

c) As the last general example of canonical maps we examine the classical change of variables

$$\Omega_{\mathcal{Z}}^{(c)} : \tilde{o}(z) = \tilde{o}(T(z_1), z_2/T'(z_1)) \quad (33)$$

or $1 \leftrightarrow 2$ on both sides. Here $T(z_1)$ describes an invertible and $\hbar$ -independent function. Using (33) in (3) one finds

$$\begin{aligned} o'(y, x) &= T'(x_+) \\ &\times o(T(x_+) + \frac{x_-}{2} T'(x_+), T(x_+) - \frac{x_-}{2} T'(x_+)) \end{aligned} \quad (34)$$

This example is also manifestly nonseparable for general cases.

These two type of maps in (b) and (c) can transform a local (separable) picture into a nonlocal (nonseparable) one.

Any nonseparable canonical map can have various compositions of elementary maps of type-II but it must also must include type-I maps such as $\Omega_{\mathcal{Z}}^{(b)}$ and/or $\Omega_{\mathcal{Z}}^{(c)}$.

Concerning the type-II case, these are standard unitary transformations of which an example we discussed as $Sp_2(\mathbb{R})$ [part (a) above]. The type-II maps within the standard Weyl formalism can be represented in the Hilbert space operator form as in (8) and the results obtained therein are valid for them. For the purpose of separability these standard maps do not offer interesting results by themselves.

5. Unitarity of the type-I maps

Coming back to type-I maps, our discussion leading to Eq. (21) classifies them also as quantum canonical maps. An interesting question is then how to incorporate them

into the quantum mechanical picture. To the author's knowledge, the type-I maps have been first examined in a quantum mechanical context by the authors of Ref. [19]. Instead of the phase space, these authors searched for the standard Hilbert space representations. The equivalence generated by the type-I maps is known as isometry, viz. they map a Hilbert space to an equivalent Hilbert space with a normally different inner product. This point has been also advocated more recently by Anderson²⁰.

In the phase space, due to their classical origin the type-I maps act on $\mathcal{Z}$. They can nevertheless be incorporated as unitary transformations in $\mathcal{Z}_*$. Therefore the function space $\mathcal{L}_2(\mathcal{Z})$ becomes the appropriate Hilbert space for such maps. That they conserve the norm of functions in $\mathcal{L}_2(\mathcal{Z})$, and hence the unitarity, can be shown by using (3) and its inverse as,

$$\begin{aligned} \int dy dx |o'(y, x)|^2 &\equiv \int \frac{d^2 z}{2\pi\hbar} \tilde{o}'^*(z) \star_z \tilde{o}'(z) \\ &= \int \frac{d^2 z}{2\pi\hbar} \tilde{o}^*(\Omega_{\mathcal{Z}} : z) \star_z \tilde{o}(\Omega_{\mathcal{Z}} : z) \\ &= \int \frac{d^2 z}{2\pi\hbar} \tilde{o}^*(Z) \star_z \tilde{o}(Z) \\ &= \int \frac{d^2 z}{2\pi\hbar} \tilde{o}^*(Z) \tilde{o}(Z) \\ &= \int \frac{d^2 Z}{2\pi\hbar} \tilde{o}^*(Z) \tilde{o}(Z) \\ &= \int \frac{d^2 Z}{2\pi\hbar} \tilde{o}^*(Z) \star_Z \tilde{o}(Z) \\ &\equiv \int dy dx |o(y, x)|^2. \end{aligned} \quad (35)$$

The preservation of the norm in $\mathcal{L}_2(\mathcal{Z})$ follows by comparing the top and the second line from the bottom. In the first line we used $\tilde{o}'(z) = (\Omega_{\mathcal{Z}} : \tilde{o})(z)$. The second and the third lines are the application of classical covariance in (13). The fourth line is the general property of the $\star$ -product, i.e. $\int d^2 z \tilde{o}_1 \star \tilde{o}_2 = \int d^2 z \tilde{o}_1 \tilde{o}_2$. In the fifth line the invariance of the integral measure, i.e. $d^2 z = d^2 Z$ is employed. The sixth line is the general property of the $\star$ -product again].

E. Nonlocality, entanglement and generalized Bell states

The key reason why a general type-I transformation is not representable locally is that such a map connects a local representation to a nonlocal one and, in this respect it can be viewed as an isometry between two Hilbert spaces which we write formally as $\hat{\Omega} : H \rightarrow H'$, where the Hilbert spaces H and H' are distinguished by their different inner products²⁰. Consider two copies H_x and H_y of the same Hilbert space and an operator defined in $H_x \times H_y$. In view of section D.4, a type-I

map acting on this operator entangles the local coordinates and the transformed operator does not have a direct product representation. The important point is that the same map can be represented unitarily in an appropriate two-dimensional noncommutative space. In the context of this work the noncommutative space in question is the phase space $\mathcal{Z}_*$. How $\hat{\Omega}$ induces entanglement/disentanglement in terms of the bilocal coordinates can be further demonstrated in the following example. Consider for instance the operator $|n\rangle\langle m|$, where $|n\rangle$ denotes the harmonic oscillator energy eigenstates, represented bilocally in $H_x \times H_y$ or equivalently, as a generalized Wigner function¹⁴ $\tilde{W}_{n,m}(z)$. A type-I map acting on this operator can be defined in the bilocal representation by using (25) and (26) as

$$(\hat{\Omega} : |n\rangle\langle m|)(y, x) = \int du \int dv \mathcal{L}(y, x; v, u) \psi_n^*(y) \psi_m(x) \quad (36)$$

where $\psi_m(x) = \langle x|n\rangle$ is the m 'th harmonic oscillator Hermite Gaussian and

$$\mathcal{L}(y, x; v, u) = \int \frac{dz_1}{2\pi\hbar} e^{-i[Z_1(z_1, x_+)u_- - z_1x_-]} \times \delta(Z_2(z_1, x_+) - u_+) \quad (37)$$

where $x_+ = (x+y)/2$, $x_- = (x-y)/2$, $u_+(u+v)/2$, $u_- = (u-v)/2$. Equivalently, $\Omega_{\mathcal{Z}}$ acts on $\tilde{W}_{n,m}(z)$ in the phase space. Using the results in section D.1 we write this as $\Omega_{\mathcal{Z}} : \tilde{W}_{n,m}(z) = \tilde{W}_{n,m}(\Omega_{\mathcal{Z}} : z) = \tilde{W}_{n,m}(Z) = \tilde{W}'_{n,m}(z)$ where Z standardly denotes the transformed phase space variables. The $\tilde{W}_{n,m}(z)$'s form a basis (the Wigner basis) in the phase space via the orthogonality relation $\int d^2z / 2\pi\hbar \tilde{W}_{n,m}^*(z) \tilde{W}_{n',m'}(z) = \delta_{n,n'} \delta_{m,m'}$. The transformed Wigner function can be expanded in this basis as

$$\Omega_{\mathcal{Z}} : W_{n,m}(z) = \sum_{n',m'} \omega_{n',m'}^{(n,m)} \tilde{W}_{n',m'}(z) \quad (38)$$

$$\omega_{n',m'}^{(n,m)} = \int \frac{d^2z}{2\pi\hbar} \tilde{W}_{n,m}^*(z) \tilde{W}_{n',m'}(Z) .$$

Let us consider a simple case $\tilde{W}_{n,m} = \delta_{n,m} \tilde{W}_n$. From a physical point of view this represents a specific pure (number) state density matrix. The corresponding Wigner function is rotationally invariant in $\mathcal{Z}_*$ and is given by the well known expression²¹

$$\tilde{W}_n(z) = 2(-1)^n e^{-(z_1^2 + z_2^2)} L_n(2(z_1^2 + z_2^2)) . \quad (39)$$

where L_n is the n 'th Laguerre polynomial. The rotational invariance in (39) is implied by the zero eigenvalue of the *phase space angular momentum* operator $K_0 = i\hbar z_j \theta_{j,k} \partial_{z_k}$ viz., $K_0 : \tilde{W}_n(z) = 0$ [a more general fact is $K_0 : \tilde{W}_{n,m} = (n-m)\hbar \tilde{W}_{n,m}$]. Let us further assume that $[\Omega_{\mathcal{Z}}, K_0] = 0$. Then the transformed Wigner function $\tilde{W}_n(Z)$ is also rotationally invariant. It can

therefore be expanded in a rotationally invariant Wigner subbasis

$$\tilde{W}_n(Z) = \sum_{n'} \omega_{n'}^{(n)} \tilde{W}_{n'}(z) \quad (40)$$

where the normalization on both sides and the invariance of the measure $d^2z = d^2Z$ requires $\sum_{n'} \omega_{n'}^{(n)} = 1$. The local symmetry algebra of (40) is $sp_2(\mathbb{R})$. The non-local symmetries are generated by the remaining type-I generators forming an infinite Lie subalgebra of the canonical algebra^{20,22}. A general rotationally invariant transformation as in (40) can be generated in two steps where the first step is $T_1 : (z_1, z_2) \rightarrow (J, \theta)$ with (J, θ) describing the generalized action-angle coordinates where for this example they specifically are $J = (z_1^2 + z_2^2)$, $\theta = \tan^{-1} z_1/z_2$. The second step is $T_2 : (J, \theta) \rightarrow (g(J), \theta/g'(J))$ where we require $g(J)$ to be of type-I hence it is $\hbar$ independent. The final transformation $T_2 T_1$ produces (40) and it can be written²² as a composition of the generators in the canonical Lie subgroup and $SP_2(\mathbb{R})$. Once the contact map $g(J)$ is known, the coefficients $\omega_{n'}^{(n)}$ are calculated by [from (39) and (40)]

$$\omega_{n'}^{(n)} = 2(-1)^{n+n'} \int_0^\infty dJ e^{-(J+g(J))} L_{n'}(2J) L_n(2g(J)) . \quad (41)$$

Eq.(41) characterizes a general nonlocal, radial (rotationally invariant) transformation. These transformations generate the bilocal Hilbert space analogs of the standard (entangled) Bell states as shown below. Hence they can be referred to as *generalized Bell states*. Here we continue with a specific example of a generalized Bell state. The simplest one has two nonzero coefficients in (40) as $\omega_{n'}^{(n)} = a_{n,n_1} \delta_{n',n_1} + a_{n,n_2} \delta_{n',n_2}$. Now let us find a canonical map that generates this Bell state. Consider the specific case $n = 0$ with $n_1 = 0$ and $n_2 = 2$ in (41). For $n = 0$ one of the Laguerre polynomials in (41) drops out ($L_0 = 1$). We next choose $g(x) = x - \ln[a_{0,0} + a_{0,2} L_2(2x)]$ (here we consider $a_{0,2} < a_{0,0}$ so that the logarithm is real). Using this in (41) it can be seen that

$$|0\rangle\langle 0| \rightarrow a_{0,0} |0\rangle\langle 0| + a_{0,2} |2\rangle\langle 2| \quad (42)$$

is obtained. More general polynomial or infinite series Bell states can also be obtained. For instance, consider for $n = 0$ the map $g(x) = x - \ln[\sum_{n'=0}^{M_{max}} \omega_{n'}^{(0)} L_{n'}(2x)]$ where M_{max} can be finite or infinite. This produces the map

$$|0\rangle\langle 0| \rightarrow \sum_{n'}^{M_{max}} a_{0,n'} |n'\rangle\langle n'| . \quad (43)$$

Even Laguerre polynomials are bounded from below. This insures that it is always possible to choose a well

defined (single valued and real) $g(x)$ by choosing the constant term $a_{0,0}$ appropriately and requiring the leading Laguerre function in the polynomial M_{max} to be of even order.

1. Generalized Bell states as noncommutative solitons

Recently radially symmetric nonlocal solutions similar to (40) have been observed (see for instance the first reference in Ref. [23] for a survey) in the noncommutative field theories as vacuum soliton solution in the infinite noncommutativity limit. In the context of these works $|0\rangle\langle 0|$ is an example of a level-one noncommutative soliton and Eq. (43) is an example of a unitary map between two vacuum configurations of a level-one soliton. The type-I and type-II transformations are joined in the canonical group of area preserving diffeomorphisms in the phase space. Therefore, we expected that this canonical group is related to the $U(\infty)$ symmetry²³. In the language of Ref. [23] the type-II maps are identified with the *local* whereas the type-I maps are associated with the *nonlocal* sectors in $U(\infty)$. Whether the canonical group covers this $U(\infty)$ entirely is currently under investigation. Furthermore, the type-I maps are fundamentally different from the non-unitary isometries presently discussed in the literature. For instance Harvey²⁴ studied the non-unitary *phase operator* $\hat{S} = \sum_{n=0}^{\infty} |n\rangle\langle n+1|$ in the context of noncommutative field theories as a generating map of fixed-level index noncommutative soliton solutions. The fact that $\hat{S}$ is representable in a local Hilbert space implies that its action preserves the local (nonlocal) sector; viz. $\mathcal{H}_x \times \mathcal{H}_y \rightarrow \mathcal{H}'_x \times \mathcal{H}'_y$. From the arguments above it is clear that such maps cannot create entanglement therefore they cannot induce transformations as in (43). In this view one immediate use of type-I transformations in the noncommutative field theory is in the generation of entangled vacuum soliton configurations. When the type-I maps can be used in composition with the $\hat{S}$ operator further soliton configurations can be obtained. As a typical composition of the two consider for instance the action of $\hat{S}^\dagger$ on (43) one finds a transformation from $|1\rangle\langle 1|$ to the generalized Bell state $\sum_{n'}^{M_{max}} a_{0,n'} |n'+1\rangle\langle n'+1|$ where leading term $M_{max}+1$ has an odd Laguerre polynomial.

Now we briefly consider the radially nonsymmetric configurations. We start by identifying $K_0 = i\hbar z_j \theta_{j,k} \partial_{z_k} = h_0 \star - \star h_0$ where $h_0 = (z_1^2 + z_2^2)/2$ is the harmonic oscillator Hamiltonian, as the third generator of $sp_2(\mathbb{R})$. The left and the right $\star$ multiplication is equivalent to a doubling of the degrees of freedom whence, K_0 has an infinitely degenerate eigenbasis $|n+k\rangle\langle n|$ with eigenvalue k for all $0 \leq n$. The other two generators of $sp_2(\mathbb{R})$, namely $K_\pm$ raise and lower k by unity for all n and, they are used as generators of non-unitary symmetries. These type of generators break the rotational symmetry by introducing non-zero phase space *angular momentum* k and, in the

context of Ref. [23], they generate soliton solutions with fixed angular momentum. We will examine the relations between the nonlinear canonical group and the $U(\infty)$ in this context in a separate work²⁵.

F. Discussion

The general approach to the canonical maps in general, and the type-I maps particularly, require reaching beyond the standard Hilbert space formalism. Type-I canonical maps have been so constructed can generate Darboux transformations between two partner Hamiltonians²⁶. In this context they play role in mapping one integrable system to a large set of its integrable partners. In addition to these results, the current work demonstrates that a noncommutative, nonlocal extension of the quantum mechanics can be obtained from the standard one by the use of type-I as well as type-II canonical maps in an extended phase space. Type-I maps establish unitary isomorphism in the phase space and isometry in the Hilbert space and, they characterize the nonlocal formulation of quantum mechanics. This result can be illustrated in the following figure.




FIG. 1. Schematic of the bilocal and the phase space representations in quantum mechanics and their interconnections by the type I and II canonical maps.

In the view of this figure, the standard representations are specific cases of, and can be obtained from, the bilocal ones at one end and the phase space representations at the other. The bilocal and the phase space representations are connected by a W map which is not shown there. The second case that one can obtain from the bilocal ones is the nonlocal (nonseparable in BLC) representations. These are connected to the noncommutative picture of the phase space representations by the same Weyl map that connects the general bilocal and the phase space ones. Each (local and nonlocal) representation is an independent automorphism created by the type II maps. The type I maps join these otherwise disjoint representations.

Note that, in the context of this work, the figure above resulted from a quantum mechanical analysis. One trivial

ial extension is to carry out the analysis in N coordinate ($2N$ phase space) dimensions. We have indications that for $1 < N$ the linear canonical group has nonlocal realizations²⁵. More interestingly, it has also implications for the field theories on noncommutative spaces. In particular, the field equations in such theories are reminiscent of the $\star$ -Schrödinger equation in (6) with the nonlinear field interactions added. The representations of these theories in the noncommutative space $\mathcal{Z}_\star$ as well as in the BLC can be fitted manifestly in the context of Fig.1. It is also specifically shown how nonlocal maps generate generalized Bell states. Interesting explorations of such maps exist in the generation and characterization of entangled soliton configurations in the noncommutative theories and in nonlocal quantum mechanics.

Acknowledgments

The author is thankful to C. Zachos and D. Fairlie and C. Deliduman for helpful discussions and critical comments.

-
- ¹ A. Connes, M.R. Douglas and A. Schwarz, [hep-th/9711162]
² Nathan Seiberg and Edward Witten, *String Theory and non-commutative geometry* JHEP 9909:032 (1999), [hep-th/9908142].
³ L. Mezincescu, *Star Operation in Quantum Mechanics* [hep-th/0007046]; J. Gamboa, M. Loewe and J.C. Rojas, *Non-commutative Quantum Mechanics* [hep-th/0010220]; S. Belucci, A. Nersessian and C. Sochichiu, *Two phases of the non-commutative quantum mechanics* [hep-th/0106138].
⁴ Daniela Bigatti and Leonard Susskind, *Magnetic fields, branes and noncommutative geometry* Phys. Rev. D **62**, 066004 (2000); [hep-th/9908056]
⁵ G. Dunne, R. Jackiw and C. Trunberger, Phys. Rev. D **41**, 661 (1990).
⁶ L. Susskind, *Quantum Hall Fluid and noncommutative Chern-Simons Theory*, [hep-th/0101029].
⁷ Alexios Polychronakos, *Quantum Hall States as matrix Chern-Simons theory*, [hep-th/0103013].
⁸ V.P. Nair and A.P. Polychronakos, *Quantum Mechanics on the Noncommutative Plane and Sphere* [hep-th/0011172].
⁹ M. Chaichian, A. Demichev, P. Prešnajder et al., *Quantum Theories on Noncommutative spaces with nontrivial topology: Aharonov-Bohm and Casimir effects*, [hep-th/0101209].
¹⁰ The waveoperator $\hat{\rho}$ here resembles in some sense the density operator in standard quantum mechanics but is not equal to it. Note that neither the crucial positivity condition nor the Hermiticity of the quantum mechanical density operator are imposed here and general results for $\hat{\rho}$ will be obtained without these conditions. However separable so-

- lutions $\rho = \psi_1 \psi_2$ can be considered as the matrix elements of the density operator in some nondiagonal basis. These are connected to a bona-fide Wigner function by the Weyl map in (3). We discuss this point later in Eq. (7).
¹¹ E. Gozzi and M. Reuter, I.J.M.P. **A 9**, 5801 (1994).
¹² It is not true that the doubled Hilbert space is equal to $H \times H$ but contains it instead. Nonlinear Poisson maps can transform functions in $H \times H$ onto a larger sector corresponding to the *entangled* part of the doubled Hilbert space. We thus refer to $H \times H$ as the local sector whereas the doubled Hilbert space is to be generally referred to as nonlocal. For further details see sections D and E.
¹³ A. Solov'yov, Mod. Phys. Lett. **A 15**, 2205 (2000).
¹⁴ T. Curtright, T. Uematsu and C. Zachos, J. Math. Phys. **42** 2396-2415 (2001) [hep-th/0011137].
¹⁵ The eigen problem using the $\star$ -products have been coined as the $\star$ -genvalue problems. For details, see D. Fairlie, Proc. Camb. Phil. Soc. **60**, 581 (1964); *ibid*, Mod. Phys. Lett. **A 13**, 263 (1998); T. Curtright, D. Fairlie and C. Zachos, Phys. Rev. D **58**, 025002 (1998).
¹⁶ T. Hakioglu and A. Dragt *Moyal-Lie Theory of phase space quantum mechanics* J. Phys. **A** to appear (2001).
¹⁷ Ruggero Maria Santilli, *Foundations of Theoretical Mechanics II: Birkhoffian Generalizations of Hamiltonian Mechanics* (Springer Verlag, 1983).
¹⁸ T. Hakioglu, *Lie algebraic approach in Weyl quantization: Nonlinear (canonical) maps and $\star$ -covariance*, ANL preprint ANL-HEP-PR-00-119, [quant-ph/0011076]
¹⁹ P.A. Mello and M. Moshinsky, J. Math. Phys. **16**, 2017 (1975); P. Kramer, M. Moshinsky and T.H. Seligman, J. Math. Phys. **10**, 683 (1977); J. Deenen, M. Moshinsky and T.H. Seligman, Ann. Phys. (NY) **127**, 458 (1980).
²⁰ Arlen Anderson Ann. Phys. **232**, 292 (1994).
²¹ H. Groenewold, Physica **12**, 405 (1946).
²² F. Leyvraz and T.H. Seligman, J. Math. Phys. **30**, 2512 (1989).
²³ Rajesh Gopakumar, Shiraz Minwalla and Andrew Strominger *Noncommutative Solitons* [hep-th/0003160]; *ibid* *On non-commutative multi-solitons* [hep-th/0103256].
²⁴ Jeffrey A. Harvey, *Kombaba Lectures on Noncommutative Solitons and D-Branes* [hep-th/0102076]; Jeffrey A. Harvey, Per Kraus and Finn Larsen, JHEP, **0012**, 024 (2000).
²⁵ T. Hakioglu and C. Deliduman, *Canonical transformations as generators of $U(\infty)$ on the noncommutative plane* (unpublished).
²⁶ Arlen Anderson, Phys. Rev. D **37**, 536 (1988); *ibid* Phys. Rev. **A43**, 4602 (1991).